Candidate	Centre	Candidate
Name	Number	Number
		0



GCSE

247/02

SCIENCE PHYSICS HIGHER TIER PHYSICS 3

A.M. FRIDAY, 28 May 2010 45 minutes

For E	Examiner's use	e only
Question	Maximum Mark	Mark awarded
1.	8	
2.	5	
3.	8	
4.	8	
5.	11	
6.	6	
7.	4	
Total	50	

ADDITIONAL MATERIALS

In addition to this paper you may require a calculator.

INSTRUCTIONS TO CANDIDATES

Write your name, centre number and candidate number in the spaces at the top of this page. Answer **all** questions.

Write your answers in the spaces provided in this booklet.

INFORMATION FOR CANDIDATES

The number of marks is given in brackets at the end of each question or part-question.

You are reminded of the necessity for good English and orderly presentation in your answers.

A list of equations is printed on page 2 of the examination paper. In calculations you should show all your working.

EQUATIONS

speed = gradient of a distance-time graph
distance travelled = area under a velocity-time graph
acceleration = gradient of a velocity-time graph

speed =
$$\frac{\text{distance}}{\text{time}}$$

$$v = u + at$$
 where $x = distance$

$$v^2 = u^2 + 2ax$$
 $u = initial velocity$
 $v = ut + \frac{1}{2}at^2$ $u = initial velocity$
 $v = acceleration$

$$x = ut + \frac{1}{2}at^2$$
 $a = acceleration$
 $x = \frac{1}{2}(u + v)t$ $t = time$

$$\frac{V_1}{V_2} = \frac{N_1}{N_2}$$
 where V_1 = voltage across the primary V_2 = voltage across the secondary N_1 = number of primary turns

 N_2 = number of secondary turns

 $momentum = mass \times velocity$

kinetic energy =
$$\frac{mv^2}{2}$$
, where $m = \text{mass}$, $v = \text{velocity or speed}$.

force =
$$\frac{\text{change in momentum}}{\text{time}}$$


wave speed = wavelength \times frequency

BLANK PAGE

(247-02) **Turn over.**

Answer all questions.

1. Ultrasound is used in sonar instruments to measure the depth of the sea. These instruments operate at a frequency of 50 000 Hz. The waves travel at a speed of 1500 m/s in water.



A wave is sent out from the boat to the sea bed and is received back 3 seconds later.

(a)	State why ultrasound cannot be heard by humans.	[1]
(b)	Use the equation	
	Distance = speed \times time	
	to calculate the depth of the sea.	[3]
		Depth of sea = m
(c)	State one other non-medical use of ultrasound.	[1]

(d) Use the equation

wavespeed = wavelength \times frequency

to find the wavelength of the ultrasound waves.

[3]

wavelength = m

8


(247-02) **Turn over.**

2. The "two-second rule" can be used by drivers who want to keep at a safe distance behind the car in front.



The time taken to say "ONLY A FOOL BREAKS THE TWO-SECOND RULE" is about two seconds. The car in front passes a post. You are too close if you reach the post in less than two seconds.

The graph shows the official stopping distances given in the Highway Code and the stopping distances given by the 2-second rule.




What is the stopping distance given by the Highway Code for a speed of 30 m/s? [1]
Stopping distance = m
How much further does the 2-second rule give you to stop than the Highway Code at a speed of 10 m/s? [2]
Further distance = m
Using the information from the graph, explain why the 2-second rule should not be used at speeds above 17 m/s. [1]
Explain why the Highway Code graph does not support the statement: "The stopping distance is proportional to the speed of the car". [1]

5

07

3. A mobile phone charger has a transformer in its plug. It changes a 230 V input to a 5 V output.




Give a reason for your answer.	[2]
Explain why the input voltage has to be alternating for the transformer to work.	[1]
Explain the purpose of the iron core.	[1]
Explain why an output voltage is produced.	[1]
	Explain why the input voltage has to be alternating for the transformer to work. Explain the purpose of the iron core.

(e)	Coil A has 9200 turns.
	Write down an equation as it appears on page 2 and use it to calculate the number of turns in coil B.
	Equation:
	[1]
	Calculation: [2]
	Number of turns in coil B =

(247-02) **Turn over.**

- 4. The diagram shows an uncontrolled nuclear fission reaction.
 - When a **slow-moving** neutron strikes an atom of $^{235}_{92}$ U, the atom splits.

In this reaction two **fast moving** neutrons are produced together with the radioactive fission fragments of Ba (barium) and Kr (krypton).



- (a) What name is given to an uncontrolled fission reaction?[1]
- (b) Complete the nuclear equation for this reaction. [2]

$$^{235}_{92}U + ^{1}_{0}n \longrightarrow ^{144}Ba + ^{36}Kr + 2^{1}_{0}n$$

- (c) In a nuclear reactor, the fission reaction is controlled using control rods of boron steel which readily absorb neutrons and a graphite moderator which improves the chances of uranium atoms splitting apart.
 - (i) State how the graphite moderator improves the possibility of fission of uranium. [1]


(ii) Explain how the energy released from a nuclear reactor can be increased. [1]

(ii) Explain now the energy released from a naclear reactor can be increased.

(d)	Outline the advantages of producing electricity from nuclear fusion rather than nuclear fission in the future. [3]	
		_

8

5. The graph shows the motion of a car travelling at 15 m/s colliding with a wall. The car rebounds with a speed of 5 m/s and comes to rest 2 seconds after hitting the wall. The mass of the car is 1200 kg.




(a) How can you tell from the graph that the car is hit backwards in the collision? [1]

<i>(b)</i>	(i)	Write down the duration of the collision.	Duration = s [1]
	(ii)	Calculate the change in velocity of the car	caused by the collision with the wall. [1]
			Change in velocity = m/s
	(iii)	Use the equation	
		force = $\frac{\text{change in momentum}}{\text{time}}$	
	to ca	alculate the force acting on the car during the	e collision. [2]
			Force = N
(c)		et an equation from page 2 and use it to estimate the wall.	mate the distance that the car rebounds
	Shov	w your working	
	Equa	ation	
			[1]
	Calc	ulation:	[3]
			Rebound distance = m
(d)		ne one safety feature in a modern car that re ollision and explain how it brings about that	

(a)	With the aid of a labelled diagram, describe Thomson's plum pudding model.
(b)	Thomson's plum pudding model was later replaced by the Bohr-Rutherford mowhich has the following main features:
	• At the core of every atom is a nucleus in which protons are contained, carrying
	 At the core of every atom is a nucleus in which protons are contained, carrying vast majority of the atom's mass. The electrons move in particular orbits around the nucleus.
Use	 vast majority of the atom's mass. The electrons move in particular orbits around the nucleus. The electrons cannot exist anywhere between the orbits.
	vast majority of the atom's mass. • The electrons move in particular orbits around the nucleus.
	 vast majority of the atom's mass. The electrons move in particular orbits around the nucleus. The electrons cannot exist anywhere between the orbits. this information to explain how atoms can be made to emit radiation. You should us
	 vast majority of the atom's mass. The electrons move in particular orbits around the nucleus. The electrons cannot exist anywhere between the orbits. this information to explain how atoms can be made to emit radiation. You should us
	 vast majority of the atom's mass. The electrons move in particular orbits around the nucleus. The electrons cannot exist anywhere between the orbits. this information to explain how atoms can be made to emit radiation. You should us
	 vast majority of the atom's mass. The electrons move in particular orbits around the nucleus. The electrons cannot exist anywhere between the orbits. this information to explain how atoms can be made to emit radiation. You should us
diagı	 vast majority of the atom's mass. The electrons move in particular orbits around the nucleus. The electrons cannot exist anywhere between the orbits. this information to explain how atoms can be made to emit radiation. You should us
diagı	vast majority of the atom's mass. • The electrons move in particular orbits around the nucleus. • The electrons cannot exist anywhere between the orbits. this information to explain how atoms can be made to emit radiation. You should us ram to help your answer.
diagı	vast majority of the atom's mass. • The electrons move in particular orbits around the nucleus. • The electrons cannot exist anywhere between the orbits. this information to explain how atoms can be made to emit radiation. You should us ram to help your answer.
diagı	vast majority of the atom's mass. • The electrons move in particular orbits around the nucleus. • The electrons cannot exist anywhere between the orbits. this information to explain how atoms can be made to emit radiation. You should us ram to help your answer.
diagi	vast majority of the atom's mass. • The electrons move in particular orbits around the nucleus. • The electrons cannot exist anywhere between the orbits. this information to explain how atoms can be made to emit radiation. You should us ram to help your answer.
diagi	vast majority of the atom's mass. • The electrons move in particular orbits around the nucleus. • The electrons cannot exist anywhere between the orbits. this information to explain how atoms can be made to emit radiation. You should us ram to help your answer.
diagi	vast majority of the atom's mass. • The electrons move in particular orbits around the nucleus. • The electrons cannot exist anywhere between the orbits. this information to explain how atoms can be made to emit radiation. You should us ram to help your answer.

7. The diagram shows the construction of an a.c. generator.



Explain how the following design features maximise the effectiveness of the generator. [4]

• electromagnets are used instead of permanent magnets.
• the rotating coil has many turns instead of just one.
• the electromagnets have curved pole pieces.